



Other Infections

Parents are requested to adopt the exclusion recommendations as detailed below in the interests of other children, staff and all families concerned.

MINIMUM PERIODS OF EXCLUSION FROM THE NURSERY

Other infections	Recommended period to be kept away from school, nursery or childminders	Comments
Conjunctivitis	None	If any outbreak/cluster occurs, consult the duty room
Diphtheria*	Exclusion is essential. Always consult with your local duty room	Family contacts must be excluded until cleared to return by the duty room. Preventable by vaccination. The duty room will organise any contact tracing necessary
Glandular fever	None	
Head lice	None	Treatment is recommended only in cases where live lice have been seen.
Hepatitis A*	Exclude until seven days after onset of jaundice (or even seven days after symptom onset if no jaundice)	The duty room will advise on any vaccination or other control measure that are needed for close contacts of a single case of hepatitis A and for suspected outbreaks.
Hepatitis B*,c, HIV/AIDS	None	Hepatitis B and C and HIV are bloodborne viruses that are not infectious through casual contact. For cleaning of body fluid spills. SEE: Good Hygiene Practice
Meningococcal Meningitis*/Septicaemia*	Until recovered	Some forms of meningococcal disease are preventable by vaccination (see immunisation schedule). There is no reason to exclude siblings or other close contacts of a case. In case of an outbreak, it may be necessary to provide antibiotics with or without meningococcal vaccination to close contacts. The Duty Room will advise on any action needed.
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. There is no reason to exclude siblings or other close contacts of a case. The Duty Room will give advice on any action needed.
Meningitis viral*	None	Milder illness. There is no reason to exclude siblings and other close contacts of a case. Contact tracing is not required.
MRSA	None	Good hygiene, in particular hand washing and environmental cleaning, are important to minimise any danger of spread. If further information is required, contact the Duty Room
Mumps*	Exclude child for five days after onset of swelling	Preventable by vaccination (MMR x2 doses)
Threadworms	None	Treatment is recommended for the child and household contacts
Tonsillitis	None	There are many causes, but most cases are due to viruses and do not need an antibiotic

Extract from 'Guidance on infection control in schools and other child care settings' issued by the Health Protection Agency 2016 (Tel: 02076548000).

